

PAPER NAME

**ojsadmin,+172.pdf**

AUTHOR

**Agung Permadi**

WORD COUNT

**3343 Words**

CHARACTER COUNT

**17401 Characters**

PAGE COUNT

**6 Pages**

FILE SIZE

**534.4KB**

SUBMISSION DATE

**Jul 6, 2023 3:35 PM GMT+8**

REPORT DATE

**Jul 6, 2023 3:36 PM GMT+8**

### ● 74% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 74% Internet database
- 14% Publications database
- Crossref database
- Crossref Posted Content database
- 0% Submitted Works database

### ● Excluded from Similarity Report

- Quoted material
- Cited material
- Small Matches (Less than 20 words)

# Circuit Training to Increase Cardiorespiratory Endurance in Male Basketball Players

Agung Wahyu Permadi<sup>1</sup>, I. Made Wisnu Adhi Putra<sup>2</sup>, Endang Sri Wahjuni<sup>3</sup>

<sup>1</sup>Departement of Physiotherapy, Faculty of Health, Science and Technology, University of Dhyana Pura, Badung, Bali, Indonesia. Br. Dinas Pohgending, Deaa Pitra, Penebel, Tabanan, Bali-Indonesia, 82152, <sup>2</sup>Departement of Nutrition Science, Faculty of Health, Science and Technology, University of Dhyana Pura, Badung, Bali, Indonesia. Address: Br. Dinas Dauh Pengkung, Ds. Tista. Kerambltan, Tabanan, Bali-Indonesia, 82161, <sup>3</sup>Department of Sport Science, Universitas Negeri Surabaya, Indonesia. Address: Unesa kampus Lidah, Jl Lidah Wetan Surabaya, 60213

## Abstract

In playing basketball, body fitness is important. It is closely related to biomotor abilities which consist of several components, one of which is endurance. An attack movement in a basketball game would require good cardiorespiratory endurance. Aerobic endurance is related to oxygen intake. This study aims to determine the improvement of cardiorespiratory endurance in male students who took basketball as a preferred extracurricular activity gained through circuit training. This is experimental research with one-group pretest-posttest design. The sample of this research consisted of 15 male students. Circuit training was carried out three times a week for four weeks at an exercise intensity of 65%–90% of maximum heart rate. From hypothesis testing with a paired t-test, it was found out that  $P = 0.000$  ( $0.000 < 0.05$ ), suggesting that there was a meaningful difference. This result shows that circuit training improved cardiorespiratory endurance.

**Keywords:** *Circuit training, cardiorespiratory endurance, male basketball players.*

## Introduction

Basketball games are considered to be one of the most dynamic and flexible sports which require high levels of physical fitness<sup>[1]</sup>. Physical relations are associated with biomotor abilities because biomotor abilities are the abilities to measure human performance<sup>[2]</sup>. One of these biomotor components greatly affects a person's endurance, namely resilience. Resilience is the ability of the heart, lung, and blood vessels to work optimally when carrying out activities for a long time without experiencing interference<sup>[3]</sup>. Resilience can be grouped into anaerobic resistance and aerobic resistance<sup>[4],[5]</sup>.

The training session applied by the coach was directed more to technical training and games. This affected the physical strength of poorly trained players<sup>[1],[6]</sup>.

Cardiorespiratory endurance can be increased by a variety of training techniques, one of which is circuit training<sup>[1],[7]</sup>. Circuit training is a combination of several types of exercises carried out in several training posts<sup>[2]</sup>. At each training post, an athlete will perform a predetermined type of exercise<sup>[8]</sup>. One circuit training set is said to be complete if an athlete has completed training in all training posts according to the prescribed dose. The movements included in this circuit training are as follows: push-ups, sit-ups, vertical jumps, abdominal curls, back extensions, astride jumping over benches, pull-ups, bench stepping, burpe, shuttle run, thrust squats, side bend, skipping, and running on the spot<sup>[9]</sup>.

## Corresponding Author:

**Agung Wahyu Permadi**

Departement of Physiotherapy, Faculty of Health, Science and Technology, University of Dhyana Pura, Badung, Bali, Indonesia. Br. Dinas Pohgending, Deaa Pitra, Penebel, Tabanan, Bali-Indonesia, 82152  
e-mail: agungwahyu@undhirabali.ac.id

## Material and Method

**Participants:** The population in this study was all male students who took extracurricular activities at a middle school in Denpasar, Bali. The sample in

1 This study was male students who took basketball as a preferred extracurricular activity. The sample used had to meet the following criteria: the participants were male middle school students who took basketball as an extracurricular activity, were aged 13–14 years, had low cardiorespiratory endurance of < 35 (poor), and did not take part in any cardiorespiratory resistance training program other than circuit training during the study. After each of these posts the students were given a break period of 15 to 20 seconds before proceeding to the next post. After completing one circuit, the students were also given a break period of 15 to 20 seconds.

**Circuit Training Measures:** In this study, some interviews and observations were carried out, and information related to age and some complaints experienced was generated. This would affect the daily physical activity and the training process and would let the researchers know whether the respondents observed were not too large. Several circuit training posts consisting of running on the spot, shuttle run, skipping, squats, push-ups, sit-ups for each set were established.

This exercise was performed in 2 repetitions (sets) with a break time of 15–20 seconds between stages and between circuits.

- a. Stage 1: Running on the spot. This training post lasted for 20 seconds.
- b. Stage 2: Shuttle run. This training post lasted for 30 seconds (the students run back and forth and touched the predetermined boundary line).
- c. Stage 3: Skipping or jumping rope. This training post lasted for 30 seconds (the students made a leap using the rope provided).
- d. Stage 4: Squat. This training post lasted for 30 seconds (the students stood then bent both knees to a half squatting position and repeated continuously for a specified time period).
- e. Stage 5: Push-up. This training post lasted for 30 seconds.
- f. Stage 6: Sit-up. This training post lasted for 30 seconds.




Fig. 1: Circuit training<sup>[9]</sup>

**Statistical Analyses:** This study used an experimental method with one-group pretest-post design. The data in the study were analyzed using SPSS. The analysis was conducted to describe the results of the research in the field without having to manipulate the facts. The data from the group were subjected to a Shapiro-Wilk test at a significance level of 0.05. It was used to examine the average chest expansion before and after treatment in each group.

**Finding and Results:** Table 1 shows the number of respondents based on age: 10 respondents (66.7%) were 13 years old, and the remaining 5 respondents (33.3%) were 14 years old.

**Table 1: Age distribution of respondents**

No	Age	Frequency	Percentage (%)
1	13	10	66,7
2	14	5	33,3
	<b>Total</b>	<b>15</b>	<b>100%</b>

Table 2 shows the number of respondents based on the VO<sub>2max</sub> values obtained from the Balke test before the circuit training was performed: 5 respondents (33.3%) obtained values of < 35 and fell into the very poor category, 8 respondents (53.3%) obtained values of 35–37 and fell into the poor category, and 2 other respondents (13.3%) obtained values of 38–44 and fell into the fair category.

**Table 2: Distribution of VO2 max values through Balke Test before being given Circuit Training**

The VO2 max value in the balke test	Total		
	Category	F	Percentage (%)
< 35	Very poor	5	33,3
35 – 37	Poor	8	53,3
38 – 44	Fair	2	13,3
<b>Total</b>		<b>15</b>	<b>100%</b>

Table 3 shows the number of respondents based on the VO<sub>2max</sub> values from the Balke test after the circuit training was performed: 1 respondent (6.7%) obtained a value of < 35 and fell into the very poor category, 6 respondents (40.0%) obtained values of 35–37 and fell into the poor category, and 8 respondents (53.3%) obtained values of 38–44 and fell into the fair category.

**Table 3: Distribution of VO2 max values through Balke Test after being given Circuit Training**

VO2 max value	Total		
	Category	F	Percentage (%)
< 35	Very poor	1	6,7
35 – 37	Poor	6	40,0
38 – 44	Fair	8	53,3
<b>Total</b>		<b>15</b>	<b>100%</b>

Table 4 shows that the average VO<sub>2max</sub> value obtained by a sample of 15 from the Balke test before the circuit training was performed was 35.68, the median was 36.2, the lowest value was 31.60, and the highest value was 39.60. Meanwhile, from the Balke test after the circuit training was performed to the same sample, the average VO<sub>2max</sub> value was 38, the median was 38.5, the lowest value was 33.30, and the highest value was 44.

**Table 4: Results of Measurement The average VO2 max value through the balke pre-test and post-test.**

Variable		Mean	Median	Min	Max	%
Balke test	Pre-test	35,68	36,20	31,60	39,60	6,5%
	Post-test	38,00	38,50	33,30	44,00	

Table 5 shows the results of the normality test using the Shapiro-Wilk test. The pre-test VO<sub>2max</sub> was 0.980. Because 0.980 > 0.05, the pre-test data were normally distributed. Meanwhile, the post-test VO<sub>2max</sub> was 0.848. Because 0.848 > 0.05, the post-test data were normally distributed.

**Table 5: Data Normality Test Results Measurement of VO2 max values through the balke test**

Variable		Statistics	Sig.	Interpretation
Balke test	Pre-test	0,982	0,980	Normal
	Post-test	0,969	0,848	Normal

Table 6 shows that the paired t-test comparing the pre-test and post-test VO<sub>2max</sub> values obtained from the Balke tests conducted on the sample yielded a significant result of 0.000 (0.000 < 0.05), indicating that there was a change in cardiorespiratory endurance after circuit training was performed.

**Table 6: Results of paired t-test analysis**

Results		Df	Sig	Information
Balke test	Pre-test	14	0,000	There are significant differences
	Post-test			

## Discussion

Cardiorespiratory endurance in males aged 13–14 (adolescents) can increase if training is applied in accordance with a stipulated dosage or training load<sup>[10]</sup>,<sup>[11]</sup>. Age affects all components of physical fitness, and  $VO_{2max}$  plays an important role in respiratory fitness.  $VO_{2max}$  of children aged 8–16 years shows a progressive and linear increase in peak aerobic ability. Thus, it can be increased by applying active sports such as circuit training<sup>[12]</sup>,<sup>[13]</sup>,<sup>[14]</sup>. However, circuit training produces different levels of  $VO_{2max}$ , causing non-optimal  $VO_{2max}$  achievement<sup>[6]</sup>,<sup>[15]</sup>. Circuit training is designed to develop cardiorespiratory fitness, cardiovascular endurance, flexibility, strength, and muscle endurance<sup>[16]</sup>. This exercise has a number of advantages: it can be performed in a short time period; it can be applied to one person or a group of persons; and it does not require any complicated equipment<sup>[2]</sup>. Circuit training is designed to stimulate the cardiorespiratory organs, and, as a result, the resistance aspect is emphasized<sup>[17]</sup>.

Some research studies reveal that male basketball players aged 10–12 years saw an increase of  $VO_{2max}$  after carrying out circuit training exercises for 6 weeks as well as average initial score and average final score by 7.68 ml/kg BW/minute (20.68%)<sup>[18]</sup>,<sup>[3]</sup>. According to the American College of Sports Medicine in 2006, the target heart rate range one should achieve when conducting a circuit training exercise to experience cardiorespiratory benefits is 65%–90% of the maximum heart rate<sup>[15]</sup>,<sup>[21]</sup>. This is in accordance with the results of the research conducted—that is, the dose used should be based on the size of the maximum heart rate to achieve changes in the cardiorespiratory aspect<sup>[19]</sup>. Recent studies related to exercises that have an effect on cardiorespiratory function have shown that breathing exercises are able to increase the amount of  $O_2$  intake, for instance, chest expansion, with a p value of  $<0.05$ <sup>[20]</sup>,<sup>[22]</sup>. Thus, the exercise also has an impact on the aerobic capacity of both sick patients and healthy people.

The results of this study show an increase in the cardiorespiratory endurance of male students taking basketball as an extracurricular activity based on the  $VO_{2max}$  values obtained from a Balke test ( $p=0.000$ ).

The students were given circuit training 3 times a week for 4 weeks of meetings with a training load of 65%–90% of the maximum heart rate. Each circuit training treatment consisted of 2 sets of exercises, each of which consisted of 6 types of exercises that had to be carried out in each training post provided. Based on the  $VO_{2max}$  values before the circuit training was given to the sample, 20% of the respondents fell into the very poor category, 66.7% to the poor category, and 13.3% to the fair category. The 6 types of exercises were running on the spot, shuttle run, skipping, squats, push-ups, and sit-ups. This study's results are supported by previous research that was conducted on middle school students, which reveals that 6-week circuit training exercises on leg muscle strength could increase  $VO_{2max}$ <sup>[18]</sup>,<sup>[23]</sup>,<sup>[8]</sup>.

The main limitation of our study is that we have yet to find any other types of training comparative to circuit training for increasing the cardiorespiratory fitness of middle-school basketball players, thus we are in need of literature related to other types of aerobic training. Therefore, more precisely, we recommend exercise to overcome the decline in cardiorespiratory fitness, for example, a decrease in the functional aerobic capacity.

## Conclusions

From the observations conducted three times a week for four weeks, it was found that the provision of circuit training could increase the cardiorespiratory endurance of male students who took basketball as an extracurricular activity. However, in order to gain further insights regarding the improvement of cardiorespiratory fitness of long-term male basketball players, the sample size should be greater because the fitness level of each man may vary.

**Conflict of Interest:** The authors declare that there is no conflict of interest related to this study.

**Source of Funding:** The authors declare that there is no source of funding from anyone.

**Ethical clearance:** The experiment was approved taken from by the Research Ethics Committee of Medical Faculty of Udayana University/Sanglah Hospital.

## References

1. Vasconcelos T, Hall A, Viana R. The influence of inspiratory muscle training on lung function in female basketball players - a randomized controlled trial. Porto Biomed J [Internet]. PBJ-Associação



1. Porto Biomedical/Porto Biomedical Society; 2017;(xx):10–3. Available from: <http://dx.doi.org/10.1016/j.pbj.2016.12.003>
2. Berner Y, Barer Y, Shefer G, Stern N. Circuit resistance training is an effective means to enhance muscle strength in older adults A Systematic Review and Meta-analysis. *Ageing Res Rev* [Internet]. Elsevier B.V; 2017; Available from: <http://dx.doi.org/10.1016/j.arr.2017.04.003>
3. Siahkhouhian M, Khodadadi D, Shahmoradi K. Effects of high-intensity interval training on aerobic and anaerobic indices: Comparison of physically active and inactive men. *Sci Sport* [Internet]. Elsevier Masson SAS; 2013;28(5). Available from: <http://dx.doi.org/10.1016/j.scispo.2012.11.006>
4. Meseguer Zafra M, García-Cantó E, Rodríguez García PL, Pérez-Soto JJ, Tárraga López PJ, Rosa Guillamón A, et al. Influence of a physical exercise program on VO<sub>2</sub>max in adults with cardiovascular risk factors. *Clin E Investig En Arterioscler Publ Of La Soc Esp Arterioscler* [Internet]. Sociedad Española de Arteriosclerosis; 2018;(xx). Available from: <https://ezproxy.southern.edu/login?url=http%3A%2F%2Fsearch.ebscohost.com%2Flogin.aspx%3Fdirect%3Dtrue%26db%3DCmedm%26AN%3D29395495%26site%3Dehost-live%26scope%3Dsite>
5. Getty AK, Wisdo TR, Chavis LN, Derella CC, McLaughlin KC, Perez AN, et al. Effects of circuit exercise training on vascular health and blood pressure. *Prev Med Reports* [Internet]. Elsevier; 2018;10(February):106–12. Available from: <https://doi.org/10.1016/j.pmedr.2018.02.010>
6. Broch K, Urheim S, Massey R, Stueflotten W, Fosså K, Hopp E, et al. Exercise capacity and peak oxygen consumption in asymptomatic patients with chronic aortic regurgitation. *Int J Cardiol*. 2016;223:688–92.
7. Ouergui I, Marzouki H, Houcine N, Franchini E, Gmada N, Bouhleb E. Relative and absolute reliability of specific kickboxing circuit training protocol in male kickboxers Reproductibilité relative et absolue d' un protocole de circuit. *Sci Sport* [Internet]. Elsevier Masson SAS; 2016;1-8. Available from: <http://dx.doi.org/10.1016/j.scispo.2016.01.004>
8. Bhambhani Y, Rowland G, Farag M. Effects of Circuit Training on Body Composition and Peak Cardiorespiratory Responses in Patients with Moderate to. 2005;86(February):268–76.
9. Sousa M De, Zouita A, Abderrahmane A Ben. Progressive circuit resistance training improves inflammatory biomarkers and insulin resistance in obese men. *Physiol Behav* [Internet]. Elsevier Inc; 2018;#pagerange#. Available from: <https://doi.org/10.1016/j.physbeh.2018.11.033>
10. Gontarev S, Kalac R. Association between high blood pressure, physical fitness, and fatness in adolescents. *J Phys Educ Sport*. 2016;16(2):1040-5.
11. Wu W, Yang Y, Chu I, Hsu H, Tsai F. Research in Developmental Disabilities Effectiveness of a cross-circuit exercise training program in improving the fitness of overweight or obese adolescents with intellectual disability enrolled in special education schools. *Res Dev Disabil* [Internet]. Elsevier Ltd; 2017;60:83–95. Available from: <http://dx.doi.org/10.1016/j.ridd.2016.11.005>
12. Randers MB, Hagman M, Brix J, Christensen JF, Pedersen MT, Nielsen JJ, et al. Effects of 3 months of full-court and half-court street basketball training on health profile in untrained men. *J Sport Heal Sci* [Internet]. Elsevier B.V; 2018;7(2):132–8. Available from: <https://doi.org/10.1016/j.jshs.2017.09.004>
13. Sirbu E. The effects of moderate aerobic training on cardiorespiratory parameters in healthy elderly subjects. *J Phys Educ Sport*. 2012;12(4):560–3.
14. Kato Y, Suzuki S, Uejima T, Semba H, Nagayama O, Hayama E, et al. The relationship between resting heart rate and peak VO<sub>2</sub>: A comparison of atrial fibrillation and sinus rhythm. *Eur J Prev Cardiol* [Internet]. American College of Cardiology Foundation; 2016;23(13):1429–36. Available from: <http://journals.sagepub.com/doi/10.1177/2047487316633885>
15. Gmiat A, Micielska K, Koz M, Flis DJ, Smaruj M, Kujach S, et al. Physiology & Behavior The impact of a single bout of high intensity circuit training on myokines' concentrations and cognitive functions in women of different age. 2017;179(January):290–7.
16. Romero-arenas S, Blazeovich AJ, Martínez-pascual M, Pérez-gómez J, Luque AJ, López-román FJ, et al. Effects of high-resistance circuit training in an elderly population. *EXG* [Internet]. Elsevier Inc; 2013;48(3):334–40. Available from: <http://dx.doi.org/10.1016/j.exger.2013.01.007>
17. Lehnert M, Stastny P, Sigmund M, Xaverova Z,

- 3  
 Hubnerova B, Kostrzewa M. The effect of combined machine and body weight circuit training for women on muscle strength and body composition. *J Phys Educ Sport.* 2015;15(3):561–8.
18. Plevková L, Peráčková J, Pačesová P, Kukurová K. The effects of a 6-week strength and endurance circuit training intervention on body image in Slovak primary school girls. 2018;(1):459–64.
19. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS, et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J.* 2016;37(27):2129–2200m.
20. Permadi AW. Comparison of respiratory training method for chest wall expansion in patients with chronic obstructive pulmonary disease. *J Phys Educ Sport.* 2018;18(4):2235–9.
21. Galazoulas C. Effects of static stretching duration on isokinetic peak torque in basketball players in semi-professional male basketball players. *J Phys Educ Sport.* 2016;16(2):1058–63.
22. Ocak Y, Savas S, Isik O, Ersoz Y. The Effect of Eight-week Workout Specific to Basketball on some Physical and Physiological Parameters. *Procedia - Soc Behav Sci [Internet]. Elsevier B.V;* 2014;152:1288–92. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S1877042814054317>
23. Sansone P, Tessitore A, Palauskas H, Lukonaitiene I, Tschan H, Pliauga V, et al. Physical and physiological demands and hormonal responses in basketball small-sided games with different tactical tasks and training regimes. *J Sci Med Sport [Internet]. Sports Medicine Australia;* 2018;11–5. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1440244018304638>

● **74% Overall Similarity**

Top sources found in the following databases:

- 74% Internet database
- 14% Publications database
- Crossref database
- Crossref Posted Content database
- 0% Submitted Works database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

<b>1</b>	<b>researchgate.net</b> Internet	<b>35%</b>
<b>2</b>	<b>medicopublication.com</b> Internet	<b>24%</b>
<b>3</b>	<b>ijphrd.com</b> Internet	<b>15%</b>
<b>4</b>	<b>repository.poltekeskupang.ac.id</b> Internet	<b>&lt;1%</b>