

# Comparison of the Effects of Different Physical Exercise Programs on the Functional Capacity of Patients with Heart Failure

Agung Wahyu Permadi<sup>1</sup>, Soetanto Hartono<sup>2</sup>, Endang Sri Wahjuni<sup>3</sup>

<sup>1</sup>Graduate Student, Doctoral Program in Sport Science, Universitas Negeri Surabaya, Indonesia,

<sup>2</sup>Professor, Head of Doctoral Program in Sport Sciences, Surabaya State University, Indonesia,

<sup>3</sup>Assistant Professor, Department of Sport Sciences, Universitas Negeri Surabaya, Indonesia

## Abstract

This study aims to compare the effects of Tai-Chi, treadmill, and stationary bicycle exercises on the capacity of heart function in patients with heart failure. The study was conducted at the outpatient clinic specific for Integrated Heart Service for 12 weeks. The research subjects included 40 respondents. Group 1 was given Tai-Chi exercise, group 2 was given treadmill exercise, group 3 was given stationary bicycle exercise, and group 4 was set as the control group. The finding show that all treatment groups had the same effect on the capacity of heart function ( $p < 0.05$ ). However, there was one group receiving the most influential effect compared to the other treatment groups, namely the Tai-Chi exercise group, with a value of  $p = 0.000$ . Based on the results of data analysis, it can be concluded that Tai-Chi exercise has a more significant effect on heart function capacity than treadmills and stationary bicycle exercises.

**Keywords:** functional capacity, heart failure physical exercise program, stationary bicycle, tai-Chi, treadmill

## Introduction

Heart failure has been one of the deadliest diseases impacting millions of people worldwide <sup>(1)</sup>. In Indonesia, heart failure contributes to more than 20% of population deaths in 2011. It has been known as the most dominant cause of death <sup>(2),(3)</sup>. In hospitals, the average mortality rate of patients with heart failure is very high despite being given rehabilitation and discharged from the hospitals <sup>(4)</sup>. Recent studies have shown an important factor in the recovery of heart failure, namely increasing the fitness of functional capacity, namely VO<sub>2</sub>Max or VO<sub>2</sub> peak, blood pressure, and heart rate <sup>(5)</sup>. Physiotherapists specializing in cardiovascular diseases have various effective training methods for treating cases of heart

failure that affect body function and movements <sup>(6)</sup>. One way to decrease the risk of heart failure is to do a structured and routine exercise by a physiotherapist <sup>(7),(8)</sup>. Until now, physiotherapy has collaboration with Cardiac rehabilitation exercise program after hospitalization that refers to aerobic physical exercise but there has been no research comparing the effect of that is able to compare Tai-Chi, treadmills, and stationary bicycle exercises <sup>(9)</sup>. The three exercise methods are considered very effective and optimal for measuring the impact of functional capacity such as blood pressure, pulse and VO<sub>2</sub>Max.

## Purpose of Research

In this study, we aim to evaluate the overall clinical impact of the exercises and compare them to find out which exercise program has a better impact on heart failure patients from outpatient care at hospital polyclinics in the Balinese population.

---

### Corresponding author:

**Agung Wahyu Permadi**

Email: [agungwahyu@undhirabali.ac.id](mailto:agungwahyu@undhirabali.ac.id)

ORCID ID: <https://orcid.org/0000-0002-9827-4987>, SCOPUS AUTHOR ID: 57205435109

## Methodology

### Participants

Forty patients (men = 22 and women = 18) were divided into four groups. Group 1 was given the treatment of Tai-Chi exercise (Yang style), group 2 was treated with treadmill exercise, group 3 was given stationary bicycle training and group 4 was set as the control group. All were given exercises twice a week for 12 weeks.

The patients were in the age range of 50 to 60 years with the provisions of the New York Heart Association (NYHA) class I and II, without a history of chronic comorbidities (post-stroke, diabetes mellitus and chronic lung disease). All participants were given education about the risks and benefits of the training. This study was approved by the Medical Research Ethics Commission of UNUD / Sanglah Hospital.

### Measurement

The measurement used a 6-minute walking test (6MWT). This running test was evaluated thoroughly and integrated all systems involved during exercises including the pulmonary and cardiovascular system, systemic circulation, peripheral circulation, blood, neuromuscular units, and muscle metabolism <sup>(10)</sup>.

### Statistical Analyses

The data description used the Statistical Product and Service Solution (SPSS) program for Windows. Data from the group were tested for Shapiro Wilk-Test normality with a significance level of 0.05. Data on

frequency test resulted in indicated the capacity of heart function before and after treatment of all study groups. Finally, Anova post-hoc Duncan was used to find out which variable or group has the most significant impact.

## Results

From homogeneity test (Lavene's Test), the blood pressure, pulse and VO2Max of all treatment groups showed  $p > 0.05$  indicating homogeneous data. Meanwhile, the normality test results of the total value of functional capacity before treatment and after treatment in each treatment group showed a p value greater than 0.05 ( $p > 0.05$ ), which means that the functional capacity value data before and after the treatment were normally distributed. Data which had normal distribution were then tested using the parametric test.

Table 3 shows that the Tai-Chi group had the lowest average blood pressure with a value of 114.76 and the highest was in the control group with a value of 138.65. In terms of the pulse variable, the Tai-Chi exercise group had the lowest value of 72.80 and the highest was in the control group with a value of 79.90. On the VO2Max variable, the control group showed the lowest VO2Max with a value of 25.20 and the Tai-Chi exercise combination group showed the highest VO2Max with a value of 34.70.

The average of all groups show that the Tai-Chi exercise group obtained a value of  $p = 0,000$  ( $p < 0.05$ ). This proves that the Tai-Chi exercise group had a significantly different effect than the other treatment groups on the capacity of heart function.

**Table 1. Homogeneity and Normality Test Results pre and post Treatment of All Groups**

Cardiac Functional Capacity Category	p Homogeneity Test (Levene's Test)	p Normality Test (Kolmogorov-Smirnov)
<b>pre</b>		0.308
Blood pressure	0.674	
Pulse	0.150	
VO2Max	0.537	
<b>post</b>		
Blood pressure	0.468	
Pulse	0.239	
VO2Max	0.249	

**Table 2. Heart Function Capacity Frequency Test in All Groups**

Cardiac Functional Capacity Category	Pre	Percentage (%)	Post	Percentage (%)
Blood pressure				
Optimal			20	50.0
Normal	8	16.0	14	38.0
Mild Hypertension	32	84.0	6	12.0
Moderate hypertension				
Severe hypertension				
Isolated Systolic Hypertension				
Pulse				
Very good	6	12.0	27	54.0
Well	33	86.0	12	44.0
Moderate	1	2.0	1	2.0
Less				
VO2Max				
Very Poor				
Poor	28	56.0	5	10.0
Fair	12	44.0	12	24.0
Good			20	60.0
Excellent			3	6.0
superior				

**Table 3. Anova Duncan test of group comparisons post treatment**

Group After Treatment	Variable(s)		
	Blood pressure	Pulse	VO2Max
Anova	0.000	0.000	0.000
Tai-Chi	114.76 <sup>a</sup>	72.80 <sup>a</sup>	34.70 <sup>c</sup>
Treadmill	134.33 <sup>c</sup>	77.20 <sup>b</sup>	31.70 <sup>b</sup>
Stationary bicycle	126.84 <sup>b</sup>	75.70 <sup>b</sup>	34.00 <sup>b</sup>
Control	138.65 <sup>d</sup>	79.90 <sup>c</sup>	25.20 <sup>a</sup>

**Annotation:**

1. p value <0.05 is called significantly different or significant
2. Numbers followed by the same notation belong to the same group
3. The notation “a” shows the lowest mean
4. The notation “c” and “d” indicate the highest mean

**Discussion**

Patients with heart failure tend to experience complex instability of heart function <sup>(11)</sup>. Generally, the Tai-chi, treadmill, and stationary bicycle exercises provide similar benefits to heart function <sup>(12), (13), (14)</sup>. In theory, Tai-Chi, treadmill, and stationary bicycle exercises are included in aerobic exercises used to optimize the overall capacity of heart function for heart failure patients <sup>(15),(16), (17), (18)</sup>. There is evidence explaining the difference between treadmill training with stationary bicycle exercises where both exercises have better effects on blood pressure and VO2Max <sup>(19), (20)</sup>. Tai-Chi program is effective for improving the fitness of heart function among elderly patients, given that Tai-chi is easy to practice <sup>(21), (22)</sup>. To determine the intensity of Tai-chi exercise, we measured the participants during the exercise at week 4, week 8, and week 12. Tai-chi group showed different blood pressure, pulse and vo2max responses than the treadmill and static bicycle groups. <sup>(17), (23)</sup>. In the literature, it is explained that Tai-chi can optimize blood pressure, pulse and vo2max of patients with heart failure. Recent findings show that Tai-chi experts are able to impact heart function from cardiovascular disease which suggests that Tai-chi may be an important alternative medicine for the prevention and treatment of heart failure <sup>(24)</sup>. One mechanism that explains the changes in heart function is that Tai-chi exercises Have a good effect on brain health and anxiety/ depression reduction <sup>(25), (26)</sup>.

Preliminary evidence supporting this study shows that the difference in impact when the test runs 6 minutes in a controlled clinical trial sample evaluating 100 outpatients proves that functional capacity and quality of life in patients given Tai-chi exercises experienced a greater improvement caused by stable blood pressure

and normal pulse <sup>(27)</sup>. The study also explained that according to data collected from the Framingham Heart Study and other prospective studies, most patients would die within 3 years after cardiac rehabilitation <sup>(28)</sup>. These preliminary findings suggest that future studies should assess the efficacy of tai chi interventions delivered by the community in increasing overall physical activity and the proportion of heart failure patients who achieve current recommendations for physical exercise (150 minutes of moderate intensity of aerobic physical activity per week) <sup>(29), (12), (30)</sup>. Other findings also prove that Tai-Chi exercise for 16 weeks has a significant effect of 40% on pulse and VO2Max. This is because Tai-Chi exercises maintain balanced body performance by prioritizing concentration so that the increase in VO2Max is more optimal <sup>(31)</sup>. Literature or previous research shows that Tai-chi has a high potential to be a safe and effective exercise option for optimizing blood pressure, pulse and VO2Max <sup>(32)</sup>.

Research comparing Tai-Chi, treadmills, and stationary bikes exercises has never been done. Therefore, the results of this study can only be compared with the results of previous studies comparing two exercises. Data from various studies that have been carried out still show different finding and no such research has been published in Asia, especially in Indonesia. Thus, this trial will provide an introduction of the benefit of tai chi as a potential exercise for heart failure patients who cannot or do not want to attend conventional heart rehabilitation programs such as treadmills or stationary bicycle exercises.

**Conclusion**

From the study conducted for 12 weeks, it can be concluded that among aerobic physical exercises such as Tai-Chi, treadmills, and stationary bicycles exercises, it is found that Tai-Chi exercises have the most significant impact compared to treadmills and static bicycles in optimizing functional capacity of patients with disease heart failure.

**Conflict of Interest:** The authors declare that there is no conflict of interest related to this study.

**Source of Funding :** No funding agencies were reported by the authors.

## References

1. Segiet OA, Romuk E, Nowalany-Kozielska E, Wojciechowska C, Piecuch A, Wojnicz R. The concentration of interleukin-33 in heart failure with reduced ejection fraction. *Anatol J Cardiol*. 2019;21(6):305–13.
2. Sumartono W, Sirait AM, Holy M, Thabrany H. Smoking and socio-demographic determinant of cardiovascular diseases among males 45+ years in Indonesia. *Int J Environ Res Public Health*. 2011;8(2):528–39.
3. Mao G, Cao Y, Wang B, Wang S, Chen Z, Wang J, et al. The salutary influence of forest bathing on elderly patients with chronic heart failure. *Int J Environ Res Public Health*. 2017;14(4).
4. Sinan ÜY, Ekmekçi A, Özbay B, Akçay FA, Bekar L, Koza Y, et al. The real-life data of hospitalized patients with heart failure: On behalf of the journey HF-TR study investigators. *Anatol J Cardiol*. 2019;21(1):25–30.
5. Pandey A, Kitzman DW, Brubaker P, Haykowsky MJ, Morgan T, Becton JT, et al. Response to Endurance Exercise Training in Older Adults with Heart Failure with Preserved or Reduced Ejection Fraction. *J Am Geriatr Soc*. 2017;65(8):1698–704.
6. Travensole C, Goessler K, Poton R, Pinto RR, Polito MD. Measurement of physical performance by field tests in programs of cardiac rehabilitation: a systematic review and meta-analysis. *Rev Port Cardiol [Internet]*. 2018;37(6):525–37. Available from: <http://dx.doi.org/10.1016/j.repce.2017.07.007>
7. Fleg JL. Exercise Therapy for Older Heart Failure Patients. *Heart Fail Clin [Internet]*. 2017;13(3):607–17. Available from: <http://dx.doi.org/10.1016/j.hfc.2017.02.012>
8. Dzibur EK, Poronsky CB. Exercise Therapy Benefits for Heart Failure. *J Nurse Pract [Internet]*. 2018;14(5):396–401. Available from: <https://doi.org/10.1016/j.nurpra.2018.01.019>
9. Menezes AR, Lavie CJ, Milani R V., Arena RA, Church TS. Cardiac rehabilitation and exercise therapy in the elderly: Should we invest in the aged? *J Geriatr Cardiol*. 2012;9(1):68–75.
10. Babu AS, Desai C V., Maiya AG, Guddattu V, Padmakumar R. Changes in derived measures from six-minute walk distance following home-based exercise training in congestive heart failure: A preliminary report. *Indian Heart J [Internet]*. 2016;68(4):527–8. Available from: <http://dx.doi.org/10.1016/j.ihj.2016.05.010>
11. Achttien RJ, Staal JB, van der Voort S, Kemps HM, Koers H, Jongert MWA, et al. Exercise-based cardiac rehabilitation in patients with chronic heart failure: A dutch practice guideline. *Netherlands Hear J*. 2015;23(1):6–17.
12. Zheng S, Lal S, Meier P, Sibbritt D, Zaslowski C. Protocol: The Effect of 12 Weeks of Tai Chi Practice on Anxiety in Healthy but Stressed People Compared to Exercise and Wait-list Comparison Groups: A Randomized Controlled Trial. *JAMS J Acupunct Meridian Stud [Internet]*. 2014;7(3):159–65. Available from: <http://dx.doi.org/10.1016/j.jams.2014.01.003>
13. Krishnaswami A, Ho WKW, Kwan WP, Tsou C, Rana JS, Solomon MD, et al. A pilot study to assess the utility of five established variables to standardize exercise treadmill test reporting. *Int J Cardiol [Internet]*. 2017;231:271–6. Available from: <http://dx.doi.org/10.1016/j.ijcard.2016.12.020>
14. Jehn M, Halle M, Schuster T, Hanssen H, Koehler F, Schmidt-Trucksäss A. Multivariable analysis of heart rate recovery after cycle ergometry in heart failure: Exercise in heart failure. *Hear Lung J Acute Crit Care*. 2011;40(6).
15. Salisbury DL, Whipple MO, Burt M, Brown RJL, Hirsch A, Foley C, et al. Translation of an evidence-based therapeutic exercise program for patients with peripheral artery disease. *J Vasc Nurs [Internet]*. 2018;36(1):23–33. Available from: <https://doi.org/10.1016/j.jvn.2017.09.003>
16. Vanroy C, Feys H, Swinnen A, Vanlandewijck Y, Truijien S, Vissers D, et al. Effectiveness of Active Cycling in Subacute Stroke Rehabilitation: A Randomized Controlled Trial. *Arch Phys Med Rehabil [Internet]*. 2017;98(8):1576-1585. e5. Available from: <http://dx.doi.org/10.1016/j.apmr.2017.02.004>
17. Ma C, Zhou W, Tang Q, Huang S. The impact of group-based Tai chi on health-status outcomes among community-dwelling older adults with hypertension. *Hear Lung [Internet]*. 2018;47(4):337–44. Available from: <https://doi.org/10.1016/j.hrtlng.2018.04.007>
18. Grazzi G, Mazzoni G, Myers J, Codecà L, Pasanisi G, Mandini S, et al. Determining the best percent-

- predicted equation for estimated VO<sub>2</sub> peak by a 1-km moderate perceptually-regulated treadmill walk to predict mortality in outpatients with cardiovascular disease. *J Sci Med Sport* [Internet]. 2018;21(3):307–11. Available from: <http://dx.doi.org/10.1016/j.jsams.2017.06.003>
19. Bittencourt MS, Christman MP, Hulten E, Divakaran S, Skali H, Kwong RY, et al. Comparison of the use of downstream tests after exercise treadmill testing by cardiologists versus noncardiologists. *Am J Cardiol* [Internet]. 2014;114(2):305–11. Available from: <http://dx.doi.org/10.1016/j.amjcard.2014.04.040>
  20. Forestieri P, Guizilini S, Peres M, Bublitz C, Bolzan DW, Rocco IS, et al. A cycle ergometer exercise program improves exercise capacity and inspiratory muscle function in hospitalized patients awaiting heart transplantation: A pilot study. *Brazilian J Cardiovasc Surg* [Internet]. 2016;31(5):389–95. Available from: <http://www.gnresearch.org/doi/10.5935/1678-9741.20160078>
  21. Li G, Yuan H, Zhang W. Effects of Tai Chi on health related quality of life in patients with chronic conditions: A systematic review of randomized controlled trials. *Complement Ther Med* [Internet]. 2014;22(4):743–55. Available from: <http://dx.doi.org/10.1016/j.ctim.2014.06.003>
  22. Li J, Hsu CC, Lin CT. Leisure participation behavior and psychological well-being of elderly adults: An empirical study of Tai Chi Chuan in China. *Int J Environ Res Public Health*. 2019;16(18):3387.
  23. Ren X, Li Y, Yang X, Li J, Li H, Yuan Z, et al. The effects of Tai Chi training in patients with heart failure: A systematic review and meta-analysis. *Front Physiol*. 2017;8(DEC):1–13.
  24. Zhang S, Zou L, Chen LZ, Yao Y, Loprinzi PD, Siu PM, et al. The effect of tai chi chuan on negative emotions in non-clinical populations: A meta-analysis and systematic review. *Int J Environ Res Public Health*. 2019;16(17):3033.
  25. Liu J, Li B, Shnider R. Effects of Tai Chi Training on Improving Physical Function in Patients With Coronary Heart Diseases. *J Exerc Sci Fit*. 2010;8(2):78–84.
  26. Hu YN, Chung YJ, Yu HK, Chen YC, Tsai CT, Hu GC. Effect of Tai Chi Exercise on Fall Prevention in Older Adults: Systematic Review and Meta-analysis of Randomized Controlled Trials. *Int J Gerontol* [Internet]. 2016;10(3):131–6. Available from: <http://dx.doi.org/10.1016/j.ijge.2016.06.002>
  27. Yeh GY, McCarthy EP, Wayne PM, Stevenson LW, Wood MJ, Davis RB, et al. Tai chi exercise in patients with chronic heart failure: A randomized clinical trial. *Arch Intern Med* [Internet]. 2011;171(8):750–7. Available from: <http://archinte.jamanetwork.com/article.aspx?doi=10.1001/archinternmed.2011.150>
  28. Salmoirago-Blotcher E, Wayne P, Bock BC, Dunsiger S, Wu WC, Stabile L, et al. Design and methods of the Gentle Cardiac Rehabilitation Study - A behavioral study of tai chi exercise for patients not attending cardiac rehabilitation. *Contemp Clin Trials* [Internet]. 2015;43:243–51. Available from: <http://dx.doi.org/10.1016/j.cct.2015.06.020>
  29. Huang YT, Wang CH, Wu YF. Adhering to a Tai Chi Chuan exercise program improves vascular resistance and cardiac function. *Int J Gerontol* [Internet]. 2011;5(3):150–4. Available from: <http://dx.doi.org/10.1016/j.ijge.2011.09.037>
  30. Lian Z, Yang L, Bian Y, Zeng L, Li M, Sun Y, et al. Effects of Tai chi on adults with essential hypertension in China: A systematic review and meta-analysis. *Eur J Integr Med*. 2017;12(January):153–62.
  31. Almodhy M, Ingle L, Sandercock GR. Effects of exercise-based cardiac rehabilitation on cardiorespiratory fitness: A meta-analysis of UK studies. *Int J Cardiol* [Internet]. 2016;221:644–51. Available from: <http://dx.doi.org/10.1016/j.ijcard.2016.06.101>
  32. Chan AWK, Chair SY, Lee DTF, Leung DYP, Sit JWH, Cheng HY, et al. Tai Chi exercise is more effective than brisk walking in reducing cardiovascular disease risk factors among adults with hypertension: A randomised controlled trial. *Int J Nurs Stud* [Internet]. 2018;88:44–52. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0020748918302013>

Compose

- Inbox
- Starred
- Snoozed
- Sent
- Drafts
- More

Labels +

sinargatsu@gmail.com



**International Journal of Contemporary Medicine** <editor.ijocm@gmail.com>

Apr 8, 2020, 3:34 PM



to me

I have the pleasure to inform you that your paper has been accepted for publication.

Pay 100 US\$ as manuscript handling charges.

Note: **India is in lockdown due to coronavirus. our offices are closed. we are carrying limited work from home. no promise can be made for date of publication. please contact us when the lockdown is lifted.**

You can make the payment by only this method

Bank details for Forex transactions

Name of Bank:	Dhanlaxmi Bank
Branch:	Lajpat Nagar, New Delhi
Name:	Ramesh Kumar Sharma
Account:	Current
Account Number:	017706700002220
IFSC/ RTGS code	DLXB0000177
Swift Code / BIC code	DLXBINBBDEL

Your acceptance letter would be sent after receipt of charges.

After this no further amendment will be entertained in the manuscript

**Please note that your submission is final and no further amendments would be accepted in manuscript after issue of acceptance letter. You are hereby advised again to check manuscript. Re submit file if corrections are there. New acceptance letter would not be issued to change in authors sequence or affiliation**